arXiv:1512.01474v1 [math.NT] 4 Dec 2015

MULTIPLICATIVE FUNCTIONS PRESERVING THE SUM-OF-THREE-SQUARES OPERATION

POO-SUNG PARK

Abstract. If a multiplicative function f satisfies $f(a^2+b^2+c^2) = f(a)^2 + f(b)^2 + f(c)^2$ for all positive integers a, b, and c, then f is an identity function.

1. introduction

In 1992, Spiro proved that if a multiplicative function $f : \mathbb{N} \to \mathbb{C}$ satisfies $f(p+q) = f(p) + f(q)$ for all primes p and q, then $f(n) = n$ under condition $f(p_0) \neq 0$ for some prime p_0. The term *multiplicative function* means $f(1) = 1$ and $f(mn) = f(m)f(n)$ for all coprime m and n.

Since then many mathematicians proposed similar propositions. Fang [7] showed the similar result for $f(p+q+r) = f(p) + f(q) + f(r)$ and Dubickas and Šarka [6] extended those results into $f(p_1 + p_2 + \cdots + p_k) = f(p_1) + f(p_2) + \cdots + f(p_k)$. Chung [3] classified multiplicative functions satisfying the equation $f(m^2+n^2) = f(m)^2 + f(n)^2$ for all $m, n \in \mathbb{N}$. Recently, Bašić [1] characterized all functions such that $f(m^2+n^2) = f(m)^2 + f(n)^2$, slightly different from Chung's condition. Many related probelms are found in [2, 4, 9, 5, 11, 10]. We would call these problems *Spiro problems*.

In this article, we add a new one to the list of Spiro problems by proving the following theorem.

Theorem 1.1. *If a multiplicative function f satisfies*

$$f(a^2+b^2+c^2) = f(a)^2 + f(b)^2 + f(c)^2$$

for all positive integers a, b, and c, then $f(n) = n$.

This research was supported by Basic Science Research Program through the National Research Foundation of Korea(NRF) funded by the Ministry of Education(NRF- 2013R1A1A2010614).

2. LEMMAS

First, we compute several values of $f(n)$.

Lemma 2.1. *If f satisfies the hypothesis of Theorem 1.1, we have $f(n) = n$ for $n \leq 15$ and $n = 25$.*

Proof. Since f is multiplicative, we have $f(1) = 1$. Clearly, $f(3) = f(1^2+1^2+1^2) = 3$. To find $f(2)$ let us consider

$$f(6) = f(1^2 + 1^2 + 2^2) = f(1)^2 + f(1)^2 + f(2)^2 = 2 + f(2)^2$$

Since $f(6) = f(2)f(3) = 3f(2)$, we have that $f(2) = 1$ or $f(2) = 2$ from the equation $2 + f(2)^2 = 3f(2)$. To determine which value $f(2)$ is we need more relations.

From $f(12) = f(2^2+2^2+2^2) = 3f(2)^2 = f(3)f(4) = 3f(4)$ we have $f(4) = f(2)^2$.

Furthermore, from $f(14) = f(1^2 + 2^2 + 3^2) = 10 + f(2)^2 = f(2)f(7)$ we have that $f(7) = f(2) + \frac{10}{f(2)}$.

To find the another relation between $f(2)$ and $f(7)$ consider

$$f(21) = f(1^2 + 2^2 + 4^2) = 1 + f(2)^2 + f(2)^4 = f(3)f(7) = 3f(7).$$

From the equation

$$1 + f(2) + f(2)^2 = 3\left(f(2) + \frac{10}{f(2)}\right)$$

we can conclude that $f(2) = 2$ and thus $f(4) = 4$, $f(6) = 6$, $f(7) = 7$, and $f(14) = 14$.

To find $f(5)$ consider $f(27)$ and $f(30)$. Since

$$\begin{aligned} f(27) &= f(1^2 + 1^2 + 5^2) = 2 + f(5)^2 \\ &= f(3^2 + 3^2 + 3^2) = 3f(3)^2 = 27, \end{aligned}$$

we have $f(5) = \pm 5$. On the other hand,

$$\begin{aligned} f(30) &= f(1^2 + 2^2 + 5^2) = 5 + f(5)^2 \\ &= f(5)f(6) = 6f(5). \end{aligned}$$

Thus, we have that $f(5) = 5$.

Since

$$f(50) = f(3^2 + 4^2 + 5^2) = 50$$
$$= f(2)f(25) = 2f(25),$$

we have $f(25) = 25$ and $f(50) = 50$.

Now, to find $f(8)$ consider $f(24)$. Since

$$f(24) = f(2^2 + 2^2 + 4^2) = 24$$
$$= f(3)f(8) = 3f(8),$$

we have that $f(8) = 8$.

Other $f(n)$ can be calculated as follows:

$$f(9) = f(1^2 + 2^2 + 2^2) = 9,$$
$$f(10) = f(2)f(5) = 10,$$
$$f(11) = f(1^2 + 1^2 + 3^2) = 11,$$
$$f(13) = f(1^2 + 2^2 + 3^2) = 13,$$
$$f(15) = f(3)f(5) = 15.$$

□

Before completing this section, we introduce a useful lemma, which is due to Hurwitz [8].

Lemma 2.2 (Hurwitz)**.** *The only squares that are not sums of three nonvanishing squares are the integers* 4^s *and* $25 \cdot 4^s$.

3. Proof of Theorem 1.1

Now consider $n > 15$. We use induction. Assume that $f(k) = k$ for all positive integers $k < n$.

If $n = a^2 + b^2 + c^2$ with $a, b, c \geq 1$, then

$$f(n) = f(a^2 + b^2 + c^2) = f(a)^2 + f(b)^2 + f(c)^2 = a^2 + b^2 + c^2 = n$$

since $f(a) = a, f(b) = b, f(c) = c$ by induction hypothesis.

If n cannot be represented as a sum of three nonvanishing squares, this means two cases:

(1) if $n = a^2 + b^2 + c^2$, then at least one of a, b, c should vanish, or
(2) $n \neq a^2 + b^2 + c^2$ for any $a, b, c \in \mathbb{Z}$.

Consider case (1): Suppose $n = a^2 + b^2$ with $3 \leq a \leq b$. Note that

$$5a < 2a^2 \leq a^2 + b^2 = n, \quad 3b < 4b \leq b^2 + 4 < b^2 + a^2 = n.$$

If $5 \nmid n$, then

$$\begin{aligned} f(25n) = f(25a^2 + 25b^2) &= f((5a)^2 + (3b)^2 + (4b)^2) \\ &= f(5a)^2 + f(3b)^2 + f(4b)^2 \\ &= (5a)^2 + (3b)^2 + (4b)^2 = 25n \end{aligned}$$

by the induction hypothesis. Since n is assumed to be indivisible by 5, $f(25n) = f(25)f(n) = 25f(n)$ and $f(n) = n$.

If $n = a^2 + b^2$ with $1 \leq a \leq 2$ and $4 \leq b$, then

$$5a \leq 10 < n, \quad 3b < 4b \leq b^2 < a^2 + b^2 = n$$

and thus $f(n) = n$ by the same reasoning.

It is already checked that $f(n) = n$ for $a \leq 2$ and $b \leq 3$ in Lemma 2.1.

If $n = a^2 + b^2$ is divisible by 5, let $n = 5^s k$ with $s \geq 1$ and $5 \nmid k$. We have shown that $f(5) = 5$ and $f(25) = 25$ in Lemma 2.1. If $s > 2$ is even, 5^s can be represented as a sum of three nonvanishing squares by Lemma 2.2. So $f(5^s) = 5^s$.

If $s \geq 3$ is odd, then $s - 3$ is even and

$$5^s = 5^{s-3} \times 125 = 5^{s-3}(3^2 + 4^2 + 10^2).$$

Thus, $f(5^s) = 5^s$.

We can conclude that $f(n) = f(5^s k) = f(5^s)f(k) = 5^s k = n$.

Now we should investigate the case of $n = a^2$. By Lemma 2.2 it suffices to check whether $f(4^s) = 4^s$. Note that

$$f(3 \cdot 4^s) = f((2^s)^2 + (2^s)^2 + (2^s)^2) = f(2^s)^2 + f(2^s)^2 + f(2^s)^2 = 3 \cdot 4^s$$
$$= f(3)f(4^s) = 3f(4^s).$$

Thus $f(4^s) = 4^s$ and we can conclude that $f(n) = n$ for the case (1).

Now consider the case (2): We have that such n is of the form $4^s(8t + 7)$ with $s, t \geq 0$. We may assume $n = 8t + 7$. Then, since $2(8t + 7) = 8(2t + 1) + 6$ can be represented as a sum of three squares of integers, $f(2(8t+7)) = 2(8t+7)$ and thus $f(8t + 7) = 8t + 7$ by dividing $f(2) = 2$.

From the case (1) and (2) we can conclude that $f(n) = n$.

References

[1] Bašić, B.: Characterization of arithmetic functions that preserve the sum-of-squares operation. *Acta Math. Sin.*, **30**, 689–695 (2014)

[2] Chen, K.-K., Chen, Y.-G.: On $f(p) + f(q) = f(p + q)$ for all odd primes p and q. *Publ. Math. Debrecen*, **76**, 425–430 (2010)

[3] Chung, P. V.: Multiplicative functions satisfying the equation $f(m^2+n^2) = f(m^2)+f(n^2)$. *Math. Slovaca*, **46**, 165–171 (1996)

[4] Chung, P. V., Phong, B. M.: Additive uniqueness sets for multiplicative functions. *Publ. Math. Debrecen*, **55**, 237–243 (1999)

[5] De Koninck, J.-M., Kátai, I., Phong, B. M.: A new characteristic of the identity function. *J. Number Theory*, **63**, 325–338 (1997)

[6] Dubickas, A., Šarka, P.: On multiplicative functions which are additive on sums of primes. *Aequat. Math.*, **86**, 81–89 (2013)

[7] Fang, J.-H.: A characterization of the identity function with equation $f(p+q+r) = f(p)+f(q)+f(r)$. *Combinatorica*, **31**, 697–701 (2011)

[8] Grosswald, E.: Representations of Integers as Sums of Squares. Springer, 2011

[9] Indlekofer, K.-H., Phong, B. M.: Additive uniqueness sets for multiplicative functions. *Ann. Univ. Sci. Budapest. Sect. Comput.*, **26**, 65–77 (2006)

[10] Phong, B. M.: A characterization of the identity function with an equation of Hosszú type. *Publ. Math. Debrecen*, **69**, 219–226 (2006)

[11] Phong, B. M.: On sets characterizing the identity function. Ann. Univ. Sci. Budapest. Sect. Comput., **24**, 295–306 (2004)

[12] Spiro, C. A.: Additive uniqueness sets for arithmetic functions. *J. Number Theory*, 42, 232–246 (1992)

Department of Mathematics Education, Kyungnam University, Changwon, Republic of Korea

E-mail address: pspark@kyungnam.ac.kr